
Suicide Decedents in Virginia: Current or Former Service Members 2003-2017

Alexandra Jansson, MPP
Program Manager, Virginia Violent Death Reporting System
Governor's Challenge - July 10, 2019
Richmond, VA



Virginia Violent Death Reporting System



- Surveil all suicides, homicides, legal interventions, deaths of undetermined intent, unintentional firearms, deaths due to terrorism
- **OCME**
- Data Sources
 - Required: medical examiner, death certificate, law enforcement, and toxicology
 - Supplemental sources

States added by year:

- 2002: MA, MD, NJ, OR, SC, VA
- 2003: AK, CO, GA, NC, OK, RI, WI
- 2004: KY, NM, UT
- 2009: MI, OH
- 2014: AZ, CT, HI, IA, IL, IN, KS, ME, MN, NH, NY, PA, VT, WA
- 2016: AL, CA, DE, DC, LA, MO, NE, NV, Puerto Rico, WV
- 2018: AR, FL, ID, MS, MT, ND, SD, TN, TX, WY




Who is a Veteran?

- Any current or former service member
- Where reported, also track:
 - Active duty
 - Branch of Service
- Limitations:
 - Reservists are probably undercounted
 - Active duty and branch of service are not always reported
- Analysis here is for veteran and civilian
 - Civilian is those ages 18 years or older who were known or unknown to have any military service



Regions for Analysis






Number of Suicide Deaths by Veteran Status

Region	Veteran		Non-Veteran	
Region 1	643	21.1%	2,410	78.9%
Region 2	539	19.9%	2,166	80.1%
Region 3 East	348	21.2%	1,294	78.8%
Region 3 West	245	19.1%	1,041	80.9%
Region 4	481	20.3%	1,894	79.7%
Region 5	989	31.6%	2,145	68.4%
Unknown	5	17.2%	24	82.8%
TOTAL	3,250	22.8%	10,974	77.2%



Number of Veteran Suicide Deaths by Year: Virginia



Demographics and Injury

- Demographics
 - White males 80% or more in every region
 - Active Duty (~4%) is largely Navy
 - Few had been served by Veteran's Administration Medical Center (VAMC)
- Injury
 - Firearms (60%-83%)
 - Asphyxia (6%-18%)
 - Poison (5%-14%)
 - Took action to prevent suicide (23.3%)
 - 17% of these actions were to restrict access to any type of lethal means

Key Differences by Age Groups

			•
	18-34	35-64	65+
•	 While a mental health diagnosis is present for almost half of all decedents, the percentage of diagnosed decedents increases for 20-24 [39.9%] to 25-34 [53.4%]). It remains fairly steady across the remainder of the lifespan, decreasing slightly after age 65 Diagnosis - PTSD (25-34; 14.0%) Only age group where the percentage exceeds 5% Substance abuse Criminal legal problems (25-34 years) 	 Interpersonal problems Family relationship problems Intimate partner relationship problems Civil legal problems (i.e. divorce or custody cases) Alcohol abuse Mental health diagnosis - depression Current mental health treatment - highest during these ages; decreases after age 65 Job and financial problems 	 Physical health problems half or more of decedents beginning at age 65 Recent death or suicide death of a friend or family member Current depressed mood
Reactive suicides - crises, recent arguments, intimate partner problems			Less reactive suicides



2017 Highlights

- Highest number and percent of female veterans since VVDRS began collecting data (n=15)
- There were 259 violent deaths of veterans in Virginia.
 89% were suicide deaths
- Veteran decedents were 3 times as frequently diagnosed with Post Traumatic Stress Disorder (PTSD)
 - Among veterans with a mental health diagnosis, PTSD was especially elevated for ages 20-24 (37.5%), 25-34 (28%), and 35-44 (18.8%)



For more information:

Alexandra Jansson, MPP
Program Manager, Virginia Violent Death Reporting System
alexandra.jansson@vdh.virginia.gov

