



# COMMONWEALTH of VIRGINIA

## DEPARTMENT OF BEHAVIORAL HEALTH AND DEVELOPMENTAL SERVICES

Post Office Box 1797  
Richmond, Virginia 23218-1797

Nelson Smith  
Commissioner

Telephone (804) 786-3921  
Fax (804) 371-6638  
www.dbhds.virginia.gov

### Office of Integrated Health Health & Safety Alert/Information

## Stroke Awareness Health and Safety Alert

---

### What is a CVA/Stroke?

**A cerebrovascular accident (CVA), more commonly known as a stroke,** occurs when there is insufficient blood flow to a part of the brain, or when a hemorrhage occurs in the brain resulting in brain cell death (Center for Disease Control and Prevention, 2019). Stroke is the 5<sup>th</sup> leading cause of death and/or long-term disability in the United States. This equates to 795,000 individuals yearly. Nearly 160,000 people each year die due to stroke related causes. Experts estimate 80% of strokes are preventable (American Stroke Association, 2019). Focusing on the need for immediate medical care is critical in order to avoid long-term outcomes such as permanent disability or death.

### Types of Stroke

- **Ischemic** - Occurs when blood vessels supplying blood to the brain are obstructed, this accounts for 87% of all strokes (American Stroke Association, 2019).
- **Hemorrhagic** - Occurs when weakened walls of the blood vessels rupture. This type of stroke is associated with uncontrolled blood pressure.
- **TIA (Transient Ischemic Attack)** - Often referred to as a “Mini-Stroke”, are caused by temporary clots. This is a warning sign of a future more serious stroke. 15% of major strokes are preceded by a TIAs (American Stroke Association, 2019).
- **Cryptogenic** - A term that refers to a stroke without a known cause (American Stroke Association, 2019).

### Signs and Symptoms of Stroke

**A stroke is a serious medical emergency requiring immediate emergency medical attention. Call 911 and get medical help at once if individuals have one or more of the following symptoms.**

## Stroke signs and symptoms include:

- ★ **Slurred speech, difficulty repeating a simple sentence.**
- ★ **Sudden trouble seeing in one or both eyes.**
- ★ **Sudden intense headache.**
- ★ **Difficulty walking, dizziness, loss of balance and coordination.**
- ★ **Sudden confusion or difficulty being understood.**
- ★ **Facial drooping on one side of face or other, mouth droops, eye droops downward.**
- ★ **Numbness, weakness, paralysis of the face, arm, or leg especially on one side of the body.**

**It is very important to note the time symptoms start and get the individual to the Emergency Room immediately (American Stroke Association, 2019).**

## Recognition of Stroke Symptoms in IDD Individuals

Close observation is important when providing care for individuals with intellectual and developmental disabilities. Careful monitoring is especially important for those who are **non-verbal** and therefore cannot perform the speech test; those who cannot raise both arms to test weakness in order to compare one side of the body to the other; and those who cannot follow a command in order to test for changes in speech or bilateral (both sides) weakness. The following is a list of various symptoms that may indicate the individual has had a stroke:

- Changes in swallowing. This could be holding food in mouth or delayed swallowing. They may suddenly be unable to control food in their mouth or may suddenly lack the ability to keep food in mouth.
- Coughing during meals, or a sudden onset of aspiration pneumonia.
- Noted changes in facial presentation, drooping eye lid, turned down mouth, sudden onset of drooling. Tongue, jaw, or lip weakness. These changes might be indicated by food spilling over their lips.
- Weakness on one side of the body. Changes in their normal gait, dragging one foot, using the wall or furniture to steady themselves. Decreased strength in their hand that might lead to them dropping items, and/or a sudden inability to control their spoon or fork when eating.
- Balance and coordination changes. You may see a sudden increase in falls. They may be unable to perform tasks such as feeding self or holding a cup in their stroke affected hand.

- Changes in vision. The individual may not be able to report changes in vision. Watch for squinting or signs that might indicate that they are having difficulty seeing.
- Pain. The individual may not be able to report pain. Note sudden holding of head, crying, grimacing or behavior problems which might be due to their inability to communicate what they are experiencing (ASA, 2019).


## Use the F.A.S.T. Acronym for Quick and Easy Stroke Recognition

If you think someone might be having a stroke act **F. A. S. T.** and do the following simple test:

- F = Face:** Ask the person to smile.  
Does one side of the face droop?
- A = Arms:** Ask the person to raise both arms.  
Does one arm drift downward?
- S = Speech:** Ask the person to repeat a simple phrase.  
Is the speech slurred or strange?
- T = Time:** Note what time did the symptoms first appear.

**Call 911 immediately if you see any of these signs or symptoms.**

This information can help health care providers determine the best treatment for each person. Do not attempt to drive someone to the hospital. **Call an ambulance so that medical personnel can begin life-saving treatment on the way to the emergency room.**



---

## Stroke Prevention

The most effective way to decrease the risk of Stroke is PREVENTION. Managing risk factors and areas of life which can be modified produces positive outcomes (NINDS, 2019).

### Non-Modifiable Risk Factors (*Things We Cannot Change*)

#### ➤ Age, Ethnicity/Race, Gender and Family History

Non-modifiable risk factors are risks none of us has any control over. African Americans have 2 times the risk for a stroke than Caucasians. If a parent has suffered a stroke before the age of 65, a person has 3 times increased risk for having a stroke than the average person (ASA, 2019).

#### ➤ Chronic Medical Conditions

People living with lifelong disabilities are more likely to have chronic medical conditions (Johnson, 2012), and the degree of disability increases the risk of poor health when compared to people with no disability (Ibarra, 2012). The following genetic conditions can increase an individual's risk for stroke for various reasons: Cerebral Autosomal Dominant Arteriopathy, Cerebral Autosomal Recessive Arteriopathy, Cerebral Amyloid Angiopathy, Moyamoya Syndrome, Fabry Disease, Ehlers-Danlos Syndrome type IV, Marfan Syndrome, Sneddon Syndrome, and Mitochondrial Encephalomyopathy (Arboix, 2015).

### Modifiable Risk Factors (*Things We Can Change*)

#### ➤ Regular Check-Ups

Regular medical check-ups give medical professionals a chance to review, discuss and monitor issues which impact an individual's health, such as: diet, smoking, weight and blood pressure (ASA, 2019). Caregivers and staff help individuals to remember scheduled appointment and assist them with any transportation issues they may have so they return to their physician for recommended and/or scheduled appointments. Try to be a good healthcare advocate.

#### ➤ Hypertension (High Blood Pressure)

According to the American Stroke Association 3 out of 4 people who suffer a stroke have high blood pressure, also known as hypertension (ASA, 2019). However, in the European POMONA II Study (Haveman, et al., 2011), hypertension rates in the IDD population does not show significantly higher rates than those of the general population. Those who participated in the POMONA II study had a hypertension rate of 5.8% in ages 19-34, which increased to 30.4% in those age 65 or older. History of stroke or cerebral hemorrhage was 1.5% (Haveman, et al., 2011). Another research study by Haveman, et al. (2010) entitled, "*Major health risks in aging persons with intellectual disabilities: an overview of recent studies*" revealed little to no difference in the rates of stroke prevalence between individuals with intellectual disability and those in the general population.

---

➤ **Physical Activity**

Evidence suggests adults with IDD are more likely to lead a sedentary lifestyle than the general population. In a study completed by Emerson of 1458 participants only 4% met the criteria to be labeled “physically active” and 44% or (635) of participants were deemed “physically incapable” (Emerson, 2005). In the European POMONA II Study, individuals with intellectual disabilities had low levels of physical activity and poor diets which played a major role in numerous preventable health conditions, such as Type II Diabetes, cardiovascular disease, osteoporosis, arthritis and obesity (Haveman, et al., 2011).

➤ **Obesity**

Obesity contributes to cardiovascular disease and stroke risk. Similarly, leading a sedentary lifestyle and consuming a diet high in fat increases the risks for stroke (Arboix, 2015). Having a BMI of 25.5-30 is considered overweight and a BMI greater than 30 is considered obese (Emerson, 2005). Evidence suggests health-promoting interventions which incorporate physical activity, balanced nutrition, and healthy behavior education can have a positive influence on the health and function of adults with ID (Heller, 2011). Being able to motivate activity and be a positive role model for healthy eating habits is a key role for caregivers and direct service providing staff.

➤ **Smoking**

Smoking is a risk factor for both cardiovascular disease and stroke (Arboix, 2015). Smokers have a risk for acute ischemic stroke which is almost 2 times higher when compared to non-smokers. Smoking increases the risk of thrombus formation and atherosclerotic plaque formation (heart disease) (Arboix, 2015). The European POMONA II Study by Haveman, et al. (2011) entitled, “*Aging and health status in adults with intellectual disabilities: results of the European POMONA II study*” found that smoking was more common in individuals with DD who were 65 years of age and older, than in younger adults. The study also revealed 7% of those individuals with DD who smoked, smoked more than 20 cigarettes a day.

➤ **Health Disparities**

Research reveals health disparities among older individuals with DD (65 and over) **were preventable health conditions** which had been under-diagnosed, late-diagnosed, or inadequately managed in some way (Haveman, et al., 2011). Caregivers should strive to be effective health care advocates for individuals, by doing the following:

- Alert the individual’s PCP if you notice any changes in their health status. Changes in: weight, alertness, physical endurance, vital signs, cognitive ability, speech, swallowing, eating, bowel habits, sleeping patterns, behaviors, mood, etc. can all indicate a treatable health condition which should be brought to their physician’s attention as soon as possible (Haveman et al., 2011).

---

## Existing Conditions that Could Elevate Stroke Risk *(but not limited to)*

### ➤ **Thromboembolism**

Blood clotting disorders and genetic defects affect the proteins needed for blood clotting. Individuals are at higher risk for stroke when they have family members who have had serious blood clots, the individual has a history of blood clots before age 40, and/or has had any unexplained miscarriages. Smoking, obesity, pregnancy, surgery, long trips, prolonged sitting, use of birth control and hormone therapies also are contributing factors to blood clot formation (ASA, 2019).

### ➤ **Atrial fibrillation (AF or Afib)**

Is an irregular heartbeat and increases the risk of stroke by 5 times. Atrial fib is a condition where the upper chambers of the heartbeat irregularly and cause blood to collect in the heart, possibly where a clot forms, which then travels to the brain causing a stroke (ASA, 2019).

### ➤ **Patent Foramen Oval (PFO)**

This is a hole between heart chambers, usually diagnosed at birth, which usually closes naturally by adulthood. 1 in 5 Americans have PFO's, but do not have any symptoms, many people are unaware they have the condition (ASA, 2019).

### ➤ **Large Artery Atherosclerosis**

This is a build-up of fat, cholesterol, and fibrous tissue which grows in and on artery walls causing narrowing (ASA, 2019).

**TIA's (Transient Ischemic Attacks)** - Are a serious warning sign for the risk of stroke which is often overlooked or undiagnosed. Risk for having a serious stroke may be as high as 17% within 90 days of TIA, and the greatest risk being the week after a TIA (ASA, 2019).



---

## Recommendations

### ➤ Recognition of Risk Factors and Early Warning Signs

The key to successful lowering the risk of death or disability from stroke is controlling the risk factors and recognizing the early warning signs of a stroke. The American Stroke Association (ASA, 2019) recommends the following strategies to prevent a stroke:

#### – Monitor Blood Pressure

Help individuals monitor their blood pressure on a regular basis and/or per their physician's orders. Many pharmacies have blood pressure monitoring stations available at no charge to check blood pressure. It is recommended individuals with a diagnosis of hypertension have their blood pressure taken daily and especially daily for two weeks after treatment changes. Keeping a journal or record of readings is helpful when attending physician appointments (American Stroke Association, 2019). Home monitoring helps to obtain long-range data for the physician to evaluate, improves the assessment of hypertension drug effects, and improves compliance (Parati, 2008).

If an elevated reading occurs, wait five minutes and retake. If the reading is elevated above 180/120mm Hg and the individual is experiencing symptoms such as difficulty breathing, chest pain, weakness, back pain, difficulty speaking, or vision changes **call 911 immediately** (American Stroke Association, 2019).

In order to increase the accuracy of all blood pressure measurements, caregivers should encourage and assist the individual to remain as still and quiet as possible, while their blood pressure is being measured. Instruct individuals to sit in an upright, well-supported position with legs uncrossed. Discourage smoking, talking, eating or drinking while you are taking their blood pressure measurement. Caregivers should not take a blood pressure reading over clothing. Discourage physical activity and drinking caffeinated beverages for 30 minutes prior. It is important to take the measurement at the same time every day and at regular intervals (per physician recommendations) (American Stroke Association, 2019), (Parati, G, 2008).

Signs of elevated blood pressure are dizziness, facial flushing, and headache or vision changes. Most people do not feel any different or realize their blood pressure is elevated, thus the reason why high blood pressure (hypertension) is known as the "silent killer" (American Stroke Association, 2019).

#### – Manage Diabetes

Help and encourage individuals to keep their daily blood sugars down via diet modifications and make sure they have A1C checks on a regular basis and/or per their physician's orders. Consider a consultation with a nutritionist for meal planning (ASA, 2019).

---

### – Physical Activity

Help individuals increase their physical activity, if cleared with their physician. The recommendation for exercise for the general public is 30 minutes of exercise daily. Check with the individual's primary care physician prior to initiating **any** exercise program for **any** individual to ensure their health and safety (ASA, 2019).

### – Healthy Diet

Encourage a healthy diet to reduce cholesterol and lower BMI. Help the individual to understand the benefits of a heart healthy diet which will lower high fat consumption. Consider a consultation with a nutritionist for meal planning. Check with the individual's primary care physician for clearance and approval, prior to assisting the individual to begin any kind of weight loss or dieting program (ASA, 2019).

### – Smoking Cessation

Encourage individuals to stop smoking. Find appropriate teaching tools to help individuals understand the negative health impact of smoking. If the individual is interested in medication to help with smoking cessation, help them address the issue with their primary care physician (ASA, 2019).

### – Aspirin Therapy

Ask the individual's primary care physician if they might be a candidate for aspirin therapy to lower their stroke risk. For some individuals, aspirin therapy is not recommended. **Never** give any medication, prescribed or over the counter, to any individual without a physician's written order and instructions (ASA, 2019).

## Resources

- **American Heart Association** (<https://www.heart.org/>). Click on Health Topics to find information geared to the individual or caregiver learning experience such as healthy living, healthy eating, mental health and well-being, sleep, stress management, smoking cessation, fitness and AMA recommendations for activity. Click on Professional to get up to date information on new therapies, quality improvement, workplace health, guidelines and statements.
- **Smoking Cessation Programs for Individuals with Intellectual Disabilities**
  - The CDC (CDC, 2019b) has approved funding for all-inclusive smoking cessation programs in 19 states. You can find out more about these programs here: <https://www.cdc.gov/ncbddd/disabilityandhealth/smoking-in-adults.html>
- **National Stroke Association** (<https://www.stroke.org/understand-stroke/what-is-stroke>). This site offers detailed imagery to help explain the different types of stroke, understanding of what a stroke is, recognition of symptoms of a stroke, prevention of a stroke, the impact of a stroke, etc.



---

## References

- American Heart Association. (2019). *Heart attack and stroke symptoms*. Retrieved from <https://www.heart.org/en/about-us/heart-attack-and-stroke-symptoms>
- American Stroke Association. American Heart Association (2019). *About stroke*. Retrieved from <https://www.strokeassociation.org/en/about-stroke>
- American Stroke Association. (2019). *What are symptoms of high blood pressure*. Retrieved from <https://www.strokeassociation.org/en/health-topics/high-blood-pressure/why-high-blood-pressure-is-a-silent-killer/what-are-the-symptoms-of-high-blood-pressure>
- American Stroke Association. (2019). *Understanding blood pressure readings: Healthy and unhealthy blood pressure ranges*. Retrieved from <https://www.strokeassociation.org/en/health-topics/high-blood-pressure/understanding-blood-pressure-readings>
- American Stroke Association. (2019). *Monitoring your blood pressure at home*. Retrieved from <https://www.strokeassociation.org/en/health-topics/high-blood-pressure/understanding-blood-pressure-readings/monitoring-your-blood-pressure-at-home>
- Arboix, A. (2015). Cardiovascular risk factors for acute stroke: Risk profiles in the different subtypes of ischemic stroke. *World Journal of Clinical Cases: WJCC*, 3(5), 418.
- Centers for Disease Control and Prevention. (2019). Know the facts about stroke. Retrieved from [https://www.cdc.gov/stroke/docs/Know\\_the\\_Facts\\_About\\_Stroke.pdf](https://www.cdc.gov/stroke/docs/Know_the_Facts_About_Stroke.pdf)
- Dixon-Ibarra A, & Horner-Johnson W. (2014). Disability Status as an Antecedent to Chronic Conditions: National Health Interview Survey, 2006–2012. *Preventing Chronic Disease: Public Health Research, Practice, and Policy*, 11, 130251. DOI:10.5888/pcd11.130251.
- Emerson, E. (2005). Underweight, obesity, and exercise among adults with intellectual disabilities in supported accommodations in Northern England. *Journal of Intellectual Disability Research*, 49, p. 134-143. Doi 10.1111/j.1365-2788.2004.00617.x.
- Emerson, E., Hatton, C., Baines, S., & Robertson, J. (2016). The physical health of British adults with intellectual disability: Cross sectional study. *International Journal for Equity in Health*, 15(11). Doi: 10.1186/s12939-016-0296-x.
- Erickson, S., Kornexl, K. (2016). Blood pressure screening, control, and treatment for patients with developmental disabilities in general medicine practices. *Journal of Pharmacy Technology*, 32(6), p. 234-239. Doi 10.1177/8755-255-6663219.
- Haveman, M., Heller, T., Lee, L., Maaskant, M., Shooshtari, S., & Strydom, A. (2010). Major health risks in aging persons with intellectual disabilities: An overview of recent studies. *Journal of Policy and Practice in Intellectual Disabilities*, 7(1), p. 59-69.
- Haveman, M., Perry, J., Salvador-Carulla, L., Walsh, P., Kerr, M., Devalk, H., Hove, G., Berger, D., Azema, B., Buono, S., Cara, A., Germanavicius, A., Linehan, C., Maatta, T., Tossebro, J., & Weber, G. (2011). Aging and health status in adults with intellectual disabilities: Results of the European POMONA II study. *Journal of Intellectual & Developmental Disability*, 36(1), p. 49-60.

**DEPARTMENT OF  
BEHAVIORAL HEALTH AND DEVELOPMENTAL  
SERVICES**

Post Office Box 1797  
Richmond, Virginia 23218-1797

Telephone (804) 786-3921  
Fax (804) 371-6638  
[www.dbhds.virginia.gov](http://www.dbhds.virginia.gov)

---

Heller, T., Drum, C., McCubbin, J., & Petersen-Besse, J. (2011). Physical activity and nutrition health promotion interventions: What is working for people with intellectual disabilities? *Journal of Intellectual Developmental Disabilities*, 40(1), p. 26-36. doi:10.1352/1934-9556-49.1.26.

National Institute of Neurological Disorders and Stroke, National Institutes of Health. (2019). *Brain basics: Preventing stroke*. Retrieved from <https://www.ninds.nih.gov/health-information/public-education/brain-basics/brain-basics-preventing-stroke>

Nelson Smith  
Commissioner

**DEPARTMENT OF  
BEHAVIORAL HEALTH AND DEVELOPMENTAL  
SERVICES**

Post Office Box 1797  
Richmond, Virginia 23218-1797

Telephone (804) 786-3921  
Fax (804) 371-6638  
[www.dbhds.virginia.gov](http://www.dbhds.virginia.gov)

---